



Is lung stereotactic ablative radiotherapy safe after pneumonectomy? – a systematic review

Andrew J. Arifin¹, Faiez Al-Shafa¹, Hanbo Chen¹, R. Gabriel Boldt², Andrew Warner¹, George B. Rodrigues¹, David A. Palma¹, Alexander V. Louie²

¹Division of Radiation Oncology, London Regional Cancer Program, London, ON, Canada; ²Department of Radiation Oncology, Odette Cancer Centre, Sunnybrook Health Sciences Centre, Toronto, ON, Canada

Contributions: (I) Conception and design: AV Louie, F Al-Shafa; (II) Administrative support: AJ Arifin; (III) Provision of study materials or patients: AV Louie, RG Boldt; (IV) Collection and assembly of data: AJ Arifin, F Al-Shafa, RG Boldt; (V) Data analysis and interpretation: AJ Arifin, F Al-Shafa, A Warner, AV Louie; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Dr. Alexander V. Louie, MD, MSc, PhD. Department of Radiation Oncology, Odette Cancer Centre, Sunnybrook Health Sciences Centre, 2075 Bayview Avenue, M4N 3M5, Toronto, Ontario, Canada. Email: alexander.louie@sunnybrook.ca.

Abstract: Patients treated with surgery for lung cancer are at risk of second primary lung cancers (SPLCs), which when localized, may be amenable to radical treatment. Treatment options, however, are limited due to reduced cardiopulmonary reserve and competing mortality risks. The aim of this study was to perform a systematic review of publications examining treatment planning considerations, clinical outcomes, and toxicity rates of stereotactic ablative radiotherapy (SABR) in patients who have previously undergone pneumonectomy. A systematic review of the literature was conducted in accordance with PRISMA guidelines using PubMed and EMBASE from inception to July 2018. Articles were limited to those published in the English language. Non-review articles with patients who received exclusively lung SABR post-pneumonectomy were included. Two reviewers independently performed abstract and full-text review, with discrepancies settled by a third reviewer. Of the 215 articles identified by the initial search, 6 articles comprising 53 patients who received lung SABR post-pneumonectomy met inclusion criteria. The mean age was 68, and most patients were male (73.7%). The mean time to pneumonectomy was 6.5 years. The mean biologically effective dose was 115 Gy, and the most common dose fractionation schemes were 54 Gy in 3 fractions, 48 Gy in 4 fractions, and 50 Gy in 5 fractions. The mean follow-up was 25.4 months. The mean 1-year overall survival and 2-year local control rates were 80.6% and 89.4%. Grade 3 or higher toxicity was reported in 13.2% of patients. SABR appears to be a safe and feasible option for SPLCs in patients with prior pneumonectomy. Multi-institutional and/or prospective studies would be helpful to determine the true risk and appropriateness of SABR in this high-risk patient population.

Keywords: Stereotactic; stereotactic ablative radiotherapy (SABR); stereotactic body radiotherapy (SBRT); cancer; pneumonectomy

Submitted Aug 12, 2019. Accepted for publication Jan 02, 2020.

doi: 10.21037/tlcr.2020.01.18

View this article at: <http://dx.doi.org/10.21037/tlcr.2020.01.18>

Introduction

Lung cancer patients treated with radical intent are at a 1–2% per patient-year risk for developing a second primary lung cancer (SPLC). SPLCs can be curable, with one analysis even suggesting that its survival is potentially superior to those with initial presentation (1). Survivorship

following pneumonectomy has been referred to as having ‘a disease itself’, given the significant impact on lung function and any subsequent lung-directed treatment decision (2).

A lobectomy, the gold standard treatment for early stage non-small cell lung cancer (NSCLC), is often not feasible following pneumonectomy due to cardiopulmonary reserve




Figure 1 Flowchart of article selection.

limitations. Based on a Surveillance, Epidemiology, and End Results (SEER) analysis, sublobar resection appears to be the favored surgical approach for a SPLC post-pneumonectomy, especially when the second lesion is small (<2 cm), without locoregional spread, and where the patient has acceptable lung function (2). This, however, is not without risk, with one study reporting complication and mortality rates of 44% and 8.3% respectively (3).

In the SEER analysis, non-surgical modalities were more commonly (62.9%) employed than repeat surgery (2). Among non-surgical options, stereotactic ablative radiotherapy (SABR) is increasingly being utilized in the primary medically inoperable early stage NSCLC context, with 3-year local-regional control rates greater than 90% (4). Lung SABR is typically well-tolerated, with fatigue being the most common side effect (5). As the literature on lung SABR following pneumonectomy emerges, the aim of this study is to systematically review the literature, with a focus on treatment planning considerations, clinical outcomes, and toxicity.

Methods

A systematic review of the literature was conducted in accordance with PRISMA guidelines using the PubMed and EMBASE databases from their respective dates of inception to July 2018. A combination of search terms was designed to capture publications that reported on patients who received lung SABR (“SABR”, “SBRT”, “stereotactic”, “radiosurgery”, “SRS”, or “SRT”) after pneumonectomy.

A total of 215 unique entries were identified. Titles and abstracts were screened, and the remaining studies underwent full text review. Two reviewers independently performed abstract and full-text review, with discrepancies settled by a third reviewer. Inclusion criteria included:

- (I) Publication types: English language journal articles;
- (II) Population: prior pneumonectomy patients;
- (III) Intervention: SABR post-pneumonectomy;
- (IV) Outcome: treatment-related toxicity and cancer-related outcomes.

Articles were excluded if they were reviews, meta-analyses, guidelines or correspondences. The article selection process is illustrated in *Figure 1*. Data abstracted from the articles included study design, sample size, age, sex, follow-up duration, time from pneumonectomy, proportion of lesions biopsied, radiation planning and dosimetric parameters, toxicity, and clinical outcomes. Toxicity was graded based on Common Terminology Criteria for Adverse Events version 3.0. The weighted means were calculated by taking individual means reported from each study and weighing by their respective sample size to obtain an overall mean based on studies with available estimates. Weighted proportions were obtained using a similar approach. Biologically effective dose (BED) was calculated using the following formula:

$$BED = nd \left[1 + \frac{d}{\alpha / \beta} \right]$$

where n = number of fractions, d = dose per fraction, alpha/beta ratio = dose at which the linear and quadratic components of cell killing are equal. We assumed a tumour

Table 1 Patient characteristics of analysed studies

Characteristic	Giaj Levra <i>et al.</i>	Haasbeek <i>et al.</i>	Sainathan <i>et al.</i>	Simpson <i>et al.</i>	Testolin <i>et al.</i>	Thompson <i>et al.</i>	Weighted average or proportion
Number of patients	9	15	2	2	12	13	–
Male (%)	77.8	–	50.0	100	66.6	76.9	73.7
Median age (years)	74	–	79	69	59	69	68
COPD (%)	–	80.0	0	0	–	–	63.0
Biopsied lesions (%)	44.4	20.0	0	0	0	23.1	18.9
Peripheral lesions (%)	–	–	0	–	–	76.9	66.6
Central lesions (%)	–	–	100	–	–	23.1	33.4
Prior chest radiation (%)	–	40.0	0	0	25.0	38.5	31.8
Median time to pneumonectomy (years)	8.4	8.9	1.5	4.5	2.8	6.8	6.5

COPD, chronic obstructive pulmonary disease.

Table 2 Planning considerations of analysed studies

Characteristic	Giaj Levra <i>et al.</i>	Haasbeek <i>et al.</i>	Sainathan <i>et al.</i>	Simpson <i>et al.</i>	Testolin <i>et al.</i>	Thompson <i>et al.</i>	Weighted average or proportion
Median BED (Gy)	112.5	151.0	100.0	102.8	87.5	105.6	115
Median dose per fraction (Gy)	15	18	10	11	13.5	12	14
Median tumour diameter (mm)	24	20	–	–	–	–	22
Median GTV (cc)	–	–	–	–	2.15	4.55	3.4
Median PTV (cc)	20.9	17.0	–	–	20.7	–	19.2
3D-CRT (%)	77.8	100	0	100	0	61.5	60.4
VMAT-IGRT (%)	22.2	0	0	0	0	38.5	13.2
CyberKnife (%)	0	0	100	0	100	0	26.4
V ₅ (%)	27.9	–	–	17.5	25.2	28.6	26.7
V ₂₀ (%)	9.0	7.1	–	4.5	4.2	7.2	6.9
MLD (Gy)	6.4	–	–	3.2	–	5.6	5.7

BED, biologically effective dose; GTV, gross tumour volume; PTV, planning target volume; 3D-CRT, three-dimensional conformal radiation treatment; VMAT, volumetric modulated arc therapy; IGRT, image-guided radiation therapy; V₅, volume of lung receiving ≥ 5 Gy; V₂₀, volume of lung receiving ≥ 20 Gy; MLD, mean lung dose.

alpha/beta ratio of 10. Calculations were performed on Microsoft Excel version 16.27.

Results

Of the 215 articles identified by the initial search, 6 articles

met inclusion criteria (6-11). The majority of studies (4) were retrospective, and the remaining 2 were case reports. Publication dates ranged from 2009 to 2015.

Characteristics of the combined 53 patients identified in the studies are listed in *Table 1*. The mean age was 68, and most patients were male (73.7%). The proportion

Table 3 Outcomes of analysed studies

Outcome	Giaj Levra <i>et al.</i>	Haasbeek <i>et al.</i>	Sainathan <i>et al.</i>	Simpson <i>et al.</i>	Testolin <i>et al.</i>	Thompson <i>et al.</i>	Weighted average or proportion*
Median follow-up (months)	41.8	16.5	–	12.0	28.0	24.0	25.4
Disease-free survival (%)	3-yr: 83.0	3-yr: 80.8	–	–	2-yr: 36.1	1-yr: 82.0	–
Overall survival (%)	–	1-yr: 91.0; 2-yr: 91.0	–	–	1-yr: 80.0	1-yr: 69.0; 2-yr: 61.0	1-yr: 80.6
Local control (%)	–	2-yr: 100	–	–	2-yr: 64.5	2-yr: 100	2-yr: 89.4
Regional control (%)	–	2-yr: 80.8	–	–	–	–	–
Metastasis-free survival (%)	–	2-yr: 92.3	–	–	–	–	–
Grade 3 or higher toxicity (%)	0	13.3	100	50.0	0	15.4	13.2

*, calculated only if 3 or more studies contributed data.

of patients with chronic obstructive pulmonary disease (COPD) was 63.0%. Only one study performed pulmonary function testing prior to radiation, which reported a mean pre-SABR forced expiratory volume in 1 second (FEV₁) of 1.4 L and a mean FEV₁ change of –0.1 L after SABR (11). The proportion of lesions that were biopsied was 18.9% (range, 0–44.4%). In studies that did not biopsy all lesions, a diagnosis of SPLC was made with a combination of clinical and radiographic (including fluorodeoxyglucose-positron emission tomography) data. The majority of lesions were peripheral (weighted mean: 66.6%). The proportion of patients who received prior radiation to the chest was 31.8% (range, 0–40.0%). The mean time from pneumonectomy was 6.5 years (range, 1.5–8.9 years).

Table 2 summarizes radiation planning data. The mean BED was 115 Gy (range, 87.5–151.0 Gy), and the most common dose fractionation schemes were 54 Gy in 3 fractions, 48 Gy in 4 fractions, and 50 Gy in 5 fractions (3 of 6 studies each). All of the studies used four-dimensional computed tomography for simulation. Respiratory gating was employed in 1 patient in a single study, whereas one study used fiducials for its 12 patients. The mean tumour diameter was 22 mm and the mean gross tumour volume was 3.4 cc. The mean planning tumour volume was 19.2 cc (range, 17.0–20.9 cc). Three-dimensional conformal radiation therapy (3D-CRT) was the most common (60.4%) delivery technique, followed by CyberKnife (26.4%) and volumetric arc therapy-image guided radiation therapy (VMAT-IGRT) (13.2%). The mean lung doses were as

follows: V₅ 26.7% (range, 17.5–28.6%), V₂₀ 6.9% (range, 4.2–9.0%) and MLD 5.7 Gy (range, 3.2–6.4 Gy).

Clinical outcomes are summarized in *Table 3*. The mean follow-up was 25.4 months (range, 12–41.8 months). The mean 1-year overall survival and 2-year local control rates were 80.6% (range, 69.0–91.0%) and 89.4% (range, 64.5–100%) respectively. 3-year disease free survival rates were reported in 2 studies at 80.8% and 83.0%. One study of 15 patients reported 2-year regional control and metastasis-free survival rates of 80.8% and 92.3% respectively.

The mean rate of grade 3 or higher toxicity was 13.2%. In one study of 15 patients, 2 (13.3%) developed grade 3 radiation pneumonitis more than 4 months post-SABR (7). The two patients were noted to have higher V₂₀ and PTV compared to other patients at 10.7%/10.2% and 32.2/28.2 cc respectively, and one had prior radiation to the chest. In another study of 13 patients, 2 (15.4%) developed grade 3 radiation pneumonitis more than 3 months post-SABR (11). The V₅, V₂₀ and MLD of the patients were 43%/21%, 11%/6% and 8.2/4.6 Gy respectively, compared to 28%, 7% and 5.5 Gy in patients without radiation pneumonitis. The two patients later succumbed to cardiorespiratory failure. One case report described two cases of esophageal perforation after more than 5 months following SABR, one of which led to a patient's death (8). Both patients had central tumours (within 2 cm of main stem bronchus) treated to 50 Gy in 5 fractions every other day. Another case report described one case of death following bacterial pneumonia and uncontrollable atrial fibrillation 16 months after SABR,

though the authors comment that this is unlikely to be related to the treatment itself (9). The patient was treated with 48 Gy in 4 biweekly fractions, and had a V_5 , V_{20} and MLD of 18%, 3% and 3.0 Gy respectively. The mean rate of possible treatment-related mortality was 7.8% (4 of 51 patients).

Discussion

To our knowledge, this is the first systematic review examining clinical and planning outcomes of lung SABR in patients with previous pneumonectomy. While SABR was associated with high rates of local control, clinically significant toxicity was observed in 13.2% cases. Recognizing the limitation of the overall small sample size of patients reviewed, prior mediastinal radiation, higher lung dose and larger treatment volumes were identified as predictors for severe pneumonitis (7,11). Determining the most appropriate dose regimen and planning objectives are of specific interest, especially as SABR appeared to be highly effective with a local control rate in the mid 80 to 90% range.

Pulmonary radiofrequency ablation (RFA) may also be an option post-pneumonectomy. A recent multi-institutional retrospective study of 15 patients who had pulmonary RFA post-pneumonectomy demonstrated 95% local control, and a 2-year overall survival rate of 71.4% (12). Study patients all underwent general anesthetic and intubation. There was a single case (6.7%) of grade ≥ 3 toxicity (pneumonia requiring intravenous antibiotics), and there were no treatment-related deaths. In contrast, another multi-institutional series of 16 RFA post-pneumonectomy did not stipulate the mode of anesthesia, and reported 3 (19%) deaths treatment-related deaths, from pneumonitis (n=2) and hemothorax (repeat RFA, n=1) (13). Our study summarizes the current knowledge on the efficacy and safety of SABR in post-pneumonectomy patients and serves as a benchmark for future research. Limitations of our study include the small sample sizes that were retrospectively analyzed, and endpoint heterogeneity, limiting the generalizability of conclusions. Furthermore, an additional 5 studies that described SABR following resection were excluded from this review, as they did not stratify outcomes based on the extent of lung surgery. Ultimately, for the safety and efficacy of SABR post-pneumonectomy to be more generalizable to routine practice, we would advocate that studies comprehensively report outcomes alongside radiation planning details, so that further insight

can be gleaned on the balance between toxicity and local control in this challenging clinical scenario.

Conclusions

This systematic review demonstrates that SABR appears to be a safe and effective option for solitary pulmonary nodules in survivors of lung cancer with prior pneumonectomy. Multi-institutional and/or prospective studies would be helpful to determine the true risk and appropriateness of SABR in this high-risk patient population.

Acknowledgments

Funding: None.

Footnote

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tlcr.2020.01.18>). AV Louie has received honoraria from Varian Medical Systems Inc. and AstraZeneca. AVL serves as the unpaid editorial board member of *Translational Lung Cancer Research* from Nov 2018 to Nov 2020. The other authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Reinmuth N, Stumpf A, Stumpf P, et al. Characteristics and outcome of patients with second primary lung cancer. *Eur Respir J* 2013;42:1668-76.
2. Ayub A, Rehmani SS, Al-Ayoubi AM, et al. Pulmonary

- Resection for Second Lung Cancer After Pneumonectomy: A Population-Based Study. *Ann Thorac Surg* 2017;104:1131-7.
3. Donington JS, Miller DL, Rowland CC, et al. Subsequent pulmonary resection for bronchogenic carcinoma after pneumonectomy. *Ann Thorac Surg* 2002;74:154-8.
 4. Timmerman R, Paulus R, Galvin J, et al. Stereotactic body radiation therapy for inoperable early stage lung cancer. *JAMA* 2010;303:1070-6.
 5. Louie AV, van Werkhoven E, Chen H, et al. Patient reported outcomes following stereotactic ablative radiotherapy or surgery for stage IA non-small-cell lung cancer: Results from the ROSEL multicenter randomized trial. *Radiother Oncol* 2015;117:44-8.
 6. Gaj Levra N, Filippi AR, Guarneri A, et al. Efficacy and safety of stereotactic ablative radiotherapy in patients with previous pneumonectomy. *Tumori* 2015;101:148-53.
 7. Haasbeek CJA, Lagerwaard FJ, de Jaeger K, et al. Outcomes of stereotactic radiotherapy for a new clinical stage I lung cancer arising postpneumonectomy. *Cancer* 2009;115:587-94.
 8. Sainathan S, Wu L, Andaz S. Delayed esophageal perforation from stereotactic body radiation therapy for locally recurrent central nonsmall cell lung cancer. *Lung India* 2014;31:158.
 9. Simpson DR, Sanghvi P, Sandhu AP. Lung Stereotactic Body Radiotherapy for Early-Stage NSCLC in Patients With Prior Pneumonectomy: A Case Report. *J Natl Compr Canc Netw* 2014;12:1513-6.
 10. Testolin A, Favretto MS, Cora S, et al. Stereotactic body radiation therapy for a new lung cancer arising after pneumonectomy: dosimetric evaluation and pulmonary toxicity. *Br J Radiol* 2015;88:20150228.
 11. Thompson R, Giuliani M, Yap ML, et al. Stereotactic body radiotherapy in patients with previous pneumonectomy: safety and efficacy. *J Thorac Oncol* 2014;9:843-7.
 12. Hess A, Palussière J, Goyers JF, et al. Pulmonary radiofrequency ablation in patients with a single lung: feasibility, efficacy, and tolerance. *Radiology* 2011;258:635-42.
 13. Sofocleous CT, May B, Petre EN, et al. Pulmonary thermal ablation in patients with prior pneumonectomy. *AJR Am J Roentgenol* 2011;196:W606-12.

Cite this article as: Arifin AJ, Al-Shafa F, Chen H, Boldt RG, Warner A, Rodrigues GB, Palma DA, Louie AV. Is lung stereotactic ablative radiotherapy safe after pneumonectomy?—a systematic review. *Transl Lung Cancer Res* 2020;9(2):348-353. doi: 10.21037/tlcr.2020.01.18